

# Sharp and Simple Bounds for the raw Moments of the Binomial and Poisson Distributions

Thomas D. Ahle `thomas@ahle.dk`  
University of Copenhagen, BARC, Facebook

---

## Abstract

We prove the inequality  $E[(X/\mu)^k] \leq (\frac{k/\mu}{\log(k/\mu+1)})^k \leq \exp(k^2/(2\mu))$  for sub-Poissonian random variables, such as Binomially or Poisson distributed random variables with mean  $\mu$ . The asymptotics  $1 + O(k^2/\mu)$  can be shown to be tight for small  $k$ . This improves over previous uniform bounds for the raw moments of those distributions by a factor exponential in  $k$ .

---

## 1. Introduction

Suppose we are given an urn of  $n$  balls, each colored red with probability  $p$  and otherwise blue. What is the probability that a sample of  $k$  balls, with replacement, from this urn has only red balls? Such questions are important in sample efficient statistics and the derandomization of algorithms.

If  $X \sim \text{Binomial}(n, p)$  denotes the number of red balls in the urn, the answer is of course given by  $P = E(X/n)^k$ . A simple application of Jensen's inequality shows  $P \geq p^k$ . In the range  $k \leq \sqrt{np}$ , this is nearly tight, as previous bounds (see (1) below) were able to show  $P \leq C^k p^k$ , for some universal constant  $C > 1$ . In this note we improve the upper bounds on Binomial moments, getting rid of the  $k$  in the exponent of  $C$ . Specifically we get  $P \leq p^k e^{k^2/(2pn)}$ , which is asymptotically tight with the lower bound as  $k = o(\sqrt{np})$ .

### 1.1. Related work

A direct approach to computing the Binomial moments expands them using the Stirling numbers of the second kind [10]:  $E X^k = \sum_{i=0}^k \left\{ \begin{matrix} k \\ i \end{matrix} \right\} n^i p^i$ . This can be derived [6, 11], for example, by combining the much easier to compute factorial moments<sup>1</sup>  $E X^{\underline{k}} = n^{\underline{k}} p^k$ . Taking the first two terms,

---

<sup>1</sup>Defined  $n^{\underline{k}} = n(n-1)\cdots(n-k+1)$ .

one finds that  $\mathbb{E} X^k = (np)^k \left(1 + \binom{k}{2} \frac{1-p}{np} + O(1/n^2)\right)$  as  $n \rightarrow \infty$ . (See section 2.1 for details.) However, it is not clear what happens when  $k$  can not be considered constant with respect to  $n$ . Similarly, for the Poisson distribution the moments can be written as the so-called Bell (or Touchard) polynomials [13] in  $\mu$ :  $\mathbb{E} X^k = \sum_{i=0}^k \left\{ \begin{smallmatrix} k \\ i \end{smallmatrix} \right\} \mu^i$ . this gives a simple lower bound  $\mathbb{E} X^k \geq \mu^k \left(1 + \binom{k}{2} \frac{1}{\mu}\right)$ , matching our upper bound for  $k = O(\sqrt{\mu})$ . However the series does again not easily yield a uniform bound.

Specializing a powerful result on moments of independent random variables by Latała [7], we get for Binomial and Poisson random variables (also Pinelis [9]):

$$\left(c \frac{k/\mu}{\log(1+k/\mu)}\right)^k \leq \mathbb{E}(X/\mu)^k \leq \left(C \frac{k/\mu}{\log(1+k/\mu)}\right)^k \quad (1)$$

for some universal constants  $c < 1 < C$ . Latała's methods are general and allow for bounds of the central moments of similar strength. However, as  $k/\mu \rightarrow 0$ , the upper bound (1) is dominated by the factor  $C^k$ , which we manage to remove in this note.

Another option is the use a Rosenthal bound, such as the following by Berend and Tassa [1] (see also Johnson et al. [5])

$$\mathbb{E} X^k \leq B_k \max\{\mu, \mu^k\}. \quad (2)$$

Here  $B_k$  is the  $k$ th Bell number, which Berend and Tassa [1] show satisfies  $B_k < \left(\frac{0.792k}{\log(k+1)}\right)^k$  and it is known [2, 4] that  $B_k^{1/k} = \frac{k}{e \log k} (1 + o(1))$  as  $k \rightarrow \infty$ . The bound is unfortunately incomparable to that of this note, and of Latała, as it behaves as  $\mu$  rather than  $\mu^k$  when  $\mu < 1$ . The connection to Bell numbers is natural since we have exactly  $\mathbb{E} X^k = B_k = \sum_i \left\{ \begin{smallmatrix} k \\ i \end{smallmatrix} \right\}$  for Poisson random variables of  $\mu = 1$ .

Focusing also on Bell numbers, but using techniques similar to this note, Ostrovsky [8] gave an upper bound for Poisson moments in the case  $k \geq 2\mu$ :

$$(\mathbb{E}(X/\mu)^k)^{1/k} \leq \frac{k/\mu}{e \log(k/\mu)} \left(1 + C(\mu) \frac{\log \log(k/\mu)}{\log(k/\mu)}\right) \quad \text{if } k \geq 2\mu, \quad (3)$$

where  $C(\mu) > 0$  is some ‘‘constant’’ depending only on  $\mu$ . In the range  $k < 2\mu$ , Ostrovsky only gives the bound  $\mathbb{E}(X/\mu)^k \leq 8.9758^k$ , so similar to the bound of Latała it loses an exponential factor in  $k$  compared to Theorem 1 below.

## 2. Bounds

**Theorem 1.** *Let  $X$  be a non-negative random variable with mean  $\mu > 0$  and moment-generating function  $E[\exp(tX)]$  bounded by  $\exp(\mu(e^t - 1))$  for all  $t > 0$ . Then for all  $k > 0$ :*

$$E[X^k] \leq \left( \frac{k}{\log(1 + k/\mu)} \right)^k.$$

A standard logarithmic bound,  $\frac{x}{\log(1+x)} \leq 1 + x/2$  (see e.g. [12] eq. 6), implies the corollary

$$E[(X/\mu)^k] \leq (1 + k/(2\mu))^k \leq \exp(k^2/(2\mu)).$$

Random variables satisfying the requirement  $E[\exp(tX)] \leq \exp(\mu(e^t - 1))$  are known as sub-Poissonian and include many simple distributions, such as the Poisson or Binomial distribution. We give more examples in Section 3.

Technically our bound is shown using the moment-generating function. This in turn involves some particularly sharp bounds on the Lambert-W function, which is defined by  $W(x)e^{W(x)} = x$ .<sup>2</sup> We will use the following lemma:

**Lemma 1** (Hoorfar and Hassani [3]). *For all  $y > 1/e$  and  $x > -1/e$ ,*

$$e^{W(x)} \leq \frac{x + y}{1 + \log y}. \quad (4)$$

*Proof.* Starting from  $1 + t \leq e^t$  substitute  $\log(y) - t$  for  $t$  to get  $1 + \log y - t \leq ye^{-t}$ . Multiplying by  $e^t$  we get  $e^t(1 + \log y) \leq te^t + y$ . Let  $t = W(x)$  s.t.  $te^t = x$ . Rearranging we get (4).  $\square$

Taking  $y = e^{W(x)}$  in (4) makes the two sides equal. Hoorfar and Hassani make various substitutions, resulting in different bounds. We will be particularly interested in  $y = x + 1$ .

*Proof of Theorem 1.* Let  $m(t) = E[\exp(tX)]$  be the moment-generating function. We will bound the moments of  $X$  by

$$E[X^k] \leq m(t)(k/(et))^k, \quad (5)$$

---

<sup>2</sup>The Lambert-W function has multiple branches. We are interested in the main one, corresponding to  $x > 0$ .

which holds for all  $k \geq 0$  and  $t > 0$ . This follows from the basic inequality  $1 + x \leq e^x$ , where we substitute  $tx/k - 1$  for  $x$  to get  $tx/k \leq e^{tx/k-1} \implies x^k \leq e^{tx}(k/(et))^k$ . Taking expectations we get (5).

We now define  $B = k/\mu$  and take  $t$  such that  $te^t = B$ . Using the Lambert-W function, this means  $t = W(B)$ . We note that  $t > 0$  whenever  $B > 0$ .

We can thus bound

$$\begin{aligned}
\mathbb{E}[(X/\mu)^k] &\leq m(t)\mu^{-k}(k/(et))^k \\
&\leq \exp(\mu(e^t - 1)) \left(\frac{k}{e\mu t}\right)^k \\
&= \exp(\mu(B/t - 1)) \left(\frac{e^t}{e}\right)^k \\
&= \exp((k/B)(B/t - 1) + k(t - 1)) \\
&= \exp(kf(B)),
\end{aligned} \tag{6}$$

where  $f(B) = 1/t + t - 1 - 1/B$ .

It remains to show  $\exp(f(B)) \leq \frac{B}{\log(B+1)}$ . Taking logarithms, this means showing the bound

$$1/W(x) + W(x) - 1 - 1/x \leq \log(x) - \log \log(1 + x) \tag{7}$$

for all  $x > 0$ , where  $W(x)$  is the Lambert-W function.

We define  $g(x) = \text{LSH} - \text{RHS} = 1/W(x) + W(x) - 1 - 1/x - \log(x) + \log \log(1 + x)$ . For  $x \rightarrow 0$  we have  $1/W(x) = 1/x + 1 + O(x)$  and  $\log \log(1 + x) = \log(x) + O(x)$ . Thus  $g(x) \rightarrow 0$  as  $x \rightarrow 0$ . We proceed to show  $g(x)$  is non-increasing for  $x > 0$ , from which (7) follows.

The Lambert function is simple to differentiate using the identity  $W'(x) = \frac{W(x)}{x(1+W(x))}$ . Taking derivatives and simplifying we get  $g'(x) = \frac{1-e^{W(x)}}{x^2} + \frac{1}{(1+x)\log(1+x)}$ . Rearranging, it suffices to show the following simple bound on the Lambert function:

$$e^{W(x)} \leq \frac{x^2}{(1+x)\log(1+x)} + 1.$$

Using Lemma 1 with  $y = x + 1$  yields the bound

$$e^{W(x)} \leq \frac{2x + 1}{1 + \log(1 + x)}.$$

We then need to show  $\frac{2x+1}{1+z} \leq \frac{x^2}{(1+x)z} + 1$ , where  $z = \log(1 + x)$ . Solving the quadratic in  $z$ , we see that the inequality holds when  $x > 0$  and  $\frac{x}{1+x} \leq$

$z \leq x$ , but we have exactly that inequality for  $\log(1+x)$ , which finishes the proof.  $\square$

### 2.1. Lower bound

As mentioned in the introduction, the expansion for the Poisson moments  $\mathbb{E} X^k = \sum_{i=0}^k \left\{ \begin{smallmatrix} k \\ i \end{smallmatrix} \right\} \mu^i$  gives a simple lower bound

$$\mathbb{E} X^k \geq \mu^k \left( 1 + \frac{k(k-1)}{2\mu} \right),$$

matching theorem 1 asymptotically for  $k = O(\sqrt{\mu})$ . The expansion for Binomial moments  $\mathbb{E} X^k = \sum_{i=0}^k \left\{ \begin{smallmatrix} k \\ i \end{smallmatrix} \right\} n^i p^i$  yields a similar lower bound

$$\begin{aligned} \mathbb{E} X^k &\geq n^k p^k + \binom{k}{2} n^{k-1} p^{k-1} \\ &= (np)^k \left( \frac{n^k}{n^k} \right) \left( 1 + \binom{k}{2} \frac{1}{(n-k+1)p} \right) \\ &= (np)^k \left( \prod_{i=0}^{k-1} 1 - \frac{i}{n} \right) \left( 1 + \binom{k}{2} \frac{1}{(n-k+1)p} \right) \\ &\geq (np)^k \left( 1 - \binom{k}{2} \frac{1}{n} \right) \left( 1 + \binom{k}{2} \frac{1}{np} \right) \\ &= (np)^k \left( 1 + \binom{k}{2} \frac{1-p}{np} \left( 1 - \binom{k}{2} \frac{1}{n} \right) \right), \end{aligned}$$

which matches theorem 1 for  $k = O(\sqrt{\mu})$  and  $p$  not too close to 1.

We will investigate some more precise lower bounds as  $k/\mu$  gets large. As briefly mentioned in the introduction, there is a correspondence between the moments of a Poisson random variable and the Bell polynomials defined by  $B(k, \mu) = \sum_i \left\{ \begin{smallmatrix} k \\ i \end{smallmatrix} \right\} \mu^i$ . In particular  $\mathbb{E} X^k = B(k, \mu)$ , if  $\mu$  is the mean of the Poissonian random variable. The Bell polynomials are so named because  $B(k, 1)$  is the  $k$ th Bell number. By Dobiński's formula  $B(k, 1) = \frac{1}{e} \sum_{i=0}^{\infty} \frac{i^k}{i!}$  the Bell numbers are generalized to the real  $k$ . We write these as  $B_x = B(x, 1)$ .

We give a lower bound for  $\mathbb{E}(X/\mu)^k$  by showing the following simple connection between the Bell polynomials and Bell numbers:

**Theorem 2.** *Let  $k$  be a positive real number and  $\mu \geq 1$  be an integer. Then*

$$B(k, \mu)/\mu^k \geq B_{k/\mu}^\mu.$$

While proof below assumes  $\mu$  is an integer, we will conjecture Theorem 2 to be true for any  $\mu \geq 1$ . Now by de Bruijn's asymptotic expression for the Bell numbers [2]:

$$E(X/\mu)^k \geq B_{k/\mu}^\mu = \left( \frac{k/\mu}{e \log(k/\mu)} (1 + o(1)) \right)^k \quad \text{as } k/\mu \rightarrow \infty.$$

matching the upper bound of Ostrovsky, eq. (3), for large  $k$ , as well as Latała's uniform lower bound with a different constant.

*Proof of Theorem 2.* Let  $X, X_1, \dots, X_\mu$  be iid. Poisson variables with mean 1, then  $S = \sum_{i=1}^\mu X_i$  is Poisson with mean  $\mu$ . We write  $\|X\|_k = (E X^k)^{1/k}$ . Then by the AG inequality:

$$\|S/\mu\|_k = \left\| \frac{1}{\mu} \sum_{i=1}^\mu X_i \right\|_k \geq \left\| \left( \prod_{i=1}^\mu X_i \right)^{1/\mu} \right\|_k = \left( \prod_{i=1}^\mu \|X_i\|_k \right)^{1/\mu} = \|X\|_{k/\mu}. \quad (8)$$

Since  $X$  has mean 1 we have  $\|X\|_{k/\mu} = B_{k/\mu}^{\mu/k}$ , and as  $S$  has mean  $\mu$  we have  $\|S/\mu\|_k = B(k, \mu)^{1/k}/\mu$ . Thus eq. (8) is exactly Theorem 2.  $\square$

For small  $k/\mu$  this bound is less interesting since  $B_x \rightarrow 0$  as  $x \rightarrow 0$ , rather than 1 as our upper bound. However, it is pretty tight, as we conjecture by the following matching upper bound in terms of the Bell numbers:

**Conjecture 1.** For all  $k > 0$  and  $\mu \geq 1$ ,

$$B_{k/\mu}^{1/(k/\mu)} \leq \frac{B(k, \mu)^{1/k}}{\mu} \leq B_{k/\mu+1}^{1/(k/\mu+1)}.$$

Furthermore, for  $0 < \mu \leq 1$ ,  $\frac{B(k, \mu)^{1/k}}{\mu} \leq B_{k/\mu}^{1/(k/\mu)}$ .

While the upper bound appears true numerically, it can't follow from our moment-generating function bound eq. (6), since it drops below that for  $k/\mu$  bigger than 40. The conjectured upper bound is even incomparable with our Theorem 1, since it is slightly above  $\frac{k/\mu}{\log(1+k/\mu)}$  for very small  $k/\mu$ . In the region  $k < 2$  and  $\mu < 1$ , the conjectued bound is weaker than eq. (2) [1], but for all other parameters it is substantially tighter.

### 3. Sub-Poissonian Random Variables

We call a non-negative random variable  $X$  sub-Poissonian if  $\mathbb{E} X = \mu$  and the moment-generating function, mgf.,  $\mathbb{E} \exp(tX) \leq \exp(\mu(e^t - 1))$  for all  $t > 0$ . If  $X_1, \dots, X_n$  are sub-Poissonian with mgf.  $m_1(t), \dots, m_n(t)$ , then  $\sum_i X_i$  is sub-Poissonian as well, since

$$\mathbb{E} \exp\left(t \sum_i X_i\right) = \prod_i m_i(t) \leq \prod_i \exp(\mu_i(e^t - 1)) = \exp\left(\left(\sum_i \mu_i\right)(e^t - 1)\right).$$

A random variable bounded in  $[0, 1]$  with mean  $\mu$  has mgf.

$$\mathbb{E} \exp(tX) = 1 + \sum_{k=1}^{\infty} \frac{t^k \mathbb{E} X^k}{k!} \leq 1 + \mu \sum_{k=1}^{\infty} \frac{t^k \mathbb{E}[1^{k-1}]}{k!} = 1 + \mu(e^t - 1) \leq \exp(\mu(e^t - 1)).$$

Hence if  $X = X_1 + \dots + X_n$  where each  $X_i \in [0, 1]$  we have  $\mu = \mathbb{E} X = \sum_i \mathbb{E} X_i$  and by Theorem 1 that  $\mathbb{E}(X/\mu)^k \leq \frac{k/\mu}{\log(k/\mu+1)}$ . In particular this captures sum of Bernoulli variables with distinct probabilities.

An example of a non-sub-Poissonian distribution is the exponential distribution with mean  $1/p = \mu$ . The moment generation is  $m(t) = \frac{1}{1 - \mu(e^t - 1)}$ , which is larger than  $\exp(\mu(e^t - 1))$  for all  $t > 0$ . It is possible however that similar methods to those in the proof of Theorem 1 will still be applicable.

### 4. Acknowledgements

The author would like to thank Robert E. Gaunt for encouragement and helpful suggestions.

### References

- [1] Daniel Berend and Tamir Tassa. Improved bounds on bell numbers and on moments of sums of random variables. *Probability and Mathematical Statistics*, 30(2):185–205, 2010.
- [2] Nicolaas Govert De Bruijn. *Asymptotic methods in analysis*, volume 4. Courier Corporation, 1981.
- [3] Abdolhossein Hoorfar and Mehdi Hassani. Inequalities on the lambert w function and hyperpower function. *J. Inequal. Pure and Appl. Math*, 9(2):5–9, 2008.

- [4] Rustam Ibragimov and Sh Sharakhmetov. On an exact constant for the rosenthal inequality. *Theory of Probability & Its Applications*, 42(2):294–302, 1998.
- [5] William B Johnson, Gideon Schechtman, and Joel Zinn. Best constants in moment inequalities for linear combinations of independent and exchangeable random variables. *The Annals of Probability*, pages 234–253, 1985.
- [6] Andreas Knoblauch. Closed-form expressions for the moments of the binomial probability distribution. *SIAM Journal on Applied Mathematics*, 69(1):197–204, 2008.
- [7] Rafał Łatała et al. Estimation of moments of sums of independent real random variables. *The Annals of Probability*, 25(3):1502–1513, 1997.
- [8] E Ostrovsky and L Sirota. Non-asymptotic estimation for bell function, with probabilistic applications. *arXiv preprint arXiv:1712.08804*, 2017.
- [9] Iosif Pinelis. Optimum bounds on moments of sums of independent random vectors. *Siberian Adv. Math*, 5(3):141–150, 1995.
- [10] John Riordan. Moment recurrence relations for binomial, poisson and hypergeometric frequency distributions. *The Annals of Mathematical Statistics*, 8(2):103–111, 1937.
- [11] Maciej Skorski. Handy formulas for binomial moments. *arXiv preprint arXiv:2012.06270*, 2020.
- [12] Flemming Topsøe. Some bounds for the logarithmic function. *Inequality theory and applications*, 4(01), 2007.
- [13] Jacques Touchard. Sur les cycles des substitutions. *Acta Mathematica*, 70(1):243–297, 1939.